

Mass Formula for Self-Orthogonal Codes over $\mathbf{Z}_{p^2}$

December 23, 2008

Dedicated to Professor D. K. Ray-Chaudhuri on the occasion of his 75th birthday

ROWENA ALMA L. BETTY ¹

Graduate School of Information Sciences
Tohoku University
Sendai 980–8579, Japan

AKIHIRO MUNEMASA

Graduate School of Information Sciences
Tohoku University
Sendai 980–8579, Japan

Abstract

In this note, we establish a mass formula for self-orthogonal codes over $\mathbf{Z}_{p^2}$, where p is a prime. As a consequence, an alternative proof of the known mass formulas for self-dual codes over $\mathbf{Z}_{p^2}$ is obtained. We also establish a mass formula for even quaternary codes, which includes a mass formula for Type II quaternary codes as a special case.

¹On study leave from the Institute of Mathematics, University of the Philippines-Diliman, Quezon City 1101 Philippines

1 Introduction

In this paper, we establish a mass formula for self-orthogonal codes over $\mathbf{Z}_{p^2}$, where p is a prime. This means finding a number $M(n)$ such that

$$M(n) = \sum_{\mathcal{C}} \frac{|E|}{|\text{Aut } \mathcal{C}|}$$

where $\mathcal{C}$ runs through the equivalence classes of self-orthogonal codes of length n over $\mathbf{Z}_{p^2}$, E is the full group of all transformations that we allow in defining equivalence for code $\mathcal{C}$, and $\text{Aut } \mathcal{C}$ is the automorphism group of $\mathcal{C}$. Clearly, the mass formula gives the total number of distinct self-orthogonal codes. Mass formulas for quaternary self-dual and Type II codes were given in [6], while for odd primes p , a mass formula for self-dual codes over $\mathbf{Z}_{p^2}$ was given in [1, 2]. Mass formulas for self-dual codes over a variety of rings, taken from different references, can also be found in [10].

Unlike mass formulas for codes over finite fields, our mass formula for self-orthogonal codes over $\mathbf{Z}_{p^2}$ is given as the sum of mass formulas over some finer classes of codes. This is because codes over $\mathbf{Z}_{p^2}$ have an invariant called a type, denoted $\{k_1, k_2\}$, in place of the dimension for codes over finite fields. The type of a code over $\mathbf{Z}_{p^2}$ is determined by the dimensions of its residue and torsion. We shall determine the number of self-orthogonal codes over $\mathbf{Z}_{p^2}$ with given residue and torsion. This number is shown to be a power of the prime p , as the set of such codes has a structure of an affine subspace of the space of matrices over $\mathbf{Z}_p$. Moreover, this number depends only on the dimensions of the residue and torsion, and is independent of particular choices of a residue and torsion. When $p = 2$, there is a subclass of the class of self-orthogonal codes over $\mathbf{Z}_4$ which we call the class of even codes. When an even code over $\mathbf{Z}_4$ is self-dual, then it is called type II, which is commonly used terminology. In the literature, some authors require that type II codes over $\mathbf{Z}_4$ to contain the all-ones vector, while others do not. We shall establish mass formulas for type II codes for both variants.

The paper is organized as follows. Section 2 introduces necessary terminology. In Section 3, we consider some mappings on the space of matrices. These mappings are used in Section 4 to describe the set of generator matrices of self-orthogonal codes of given residue and torsion. In Section 5, we establish results analogous to those in Section 4, for quaternary even codes. Finally in Section 6, we give a mass formula for self-orthogonal codes

of given type, and as a corollary, we also give a mass formula for self-dual codes. Analogous results for quaternary even codes are also given.

2 Preliminaries

For a positive integer m , we denote by $\mathbf{Z}_m$ the ring of integers modulo m . A code $\mathcal{C}$ of length n over $\mathbf{Z}_m$ is a submodule of $\mathbf{Z}_m^n$. For a matrix $G \in M_{k \times n}(\mathbf{Z})$, we denote by $\mathbf{Z}_m^k G$ the code $\{aG \bmod m \mid a \in \mathbf{Z}^k\}$ of length n over $\mathbf{Z}_m$. A generator matrix of a code $\mathcal{C}$ of length n over $\mathbf{Z}_m$ is a matrix $G \in M_{k \times n}(\mathbf{Z})$ such that $\mathcal{C} = \mathbf{Z}_m^k G$. Usually, entries of a generator matrix of a code over $\mathbf{Z}_m$ are taken to be in $\mathbf{Z}_m$. However, since we deal with codes over $\mathbf{Z}_p$ and $\mathbf{Z}_{p^2}$ at the same time, we adopt this non-standard convention to avoid cumbersome notation.

We denote by $x \cdot y$ the standard inner product of vectors x, y in $\mathbf{Z}_m^n$, and by $\mathcal{C}^\perp$ the dual code of a code $\mathcal{C}$ over $\mathbf{Z}_m$ with respect to this inner product. A code $\mathcal{C}$ is said to be self-orthogonal (respectively self-dual) if $\mathcal{C} \subset \mathcal{C}^\perp$ (respectively $\mathcal{C} = \mathcal{C}^\perp$) holds. Two codes $\mathcal{C}, \mathcal{C}'$ of the same length are said to be equivalent if there exists a $(1, -1)$ monomial matrix which maps $\mathcal{C}$ onto $\mathcal{C}'$.

Let p be a prime, and consider the exact sequence

$$0 \rightarrow \mathbf{Z}_p \xrightarrow{\iota} \mathbf{Z}_{p^2} \xrightarrow{\pi} \mathbf{Z}_p \rightarrow 0,$$

where ι is the composition of the isomorphism $\mathbf{Z}_p \rightarrow p\mathbf{Z}_{p^2}$ and the embedding $p\mathbf{Z}_{p^2} \rightarrow \mathbf{Z}_{p^2}$, and π is the canonical homomorphism. For a positive integer n , by abuse of notation, we denote the cartesian product of the mappings ι and π by the same symbols:

$$0 \rightarrow \mathbf{Z}_p^n \xrightarrow{\iota} \mathbf{Z}_{p^2}^n \xrightarrow{\pi} \mathbf{Z}_p^n \rightarrow 0 \quad (\text{exact}).$$

For a code $\mathcal{C}$ over $\mathbf{Z}_{p^2}$, $\pi(\mathcal{C})$ is called the residue code of $\mathcal{C}$, and $\iota^{-1}(\mathcal{C})$ is called the torsion code of $\mathcal{C}$. Since $\iota \circ \pi = p$, we have $\pi(\mathcal{C}) \subset \iota^{-1}(\mathcal{C})$. Moreover, since $\text{Im } \iota = \text{Ker } \pi$, we have

$$|\mathcal{C}| = |\pi(\mathcal{C})| |\iota^{-1}(\mathcal{C})|. \quad (1)$$

Every code of length n over $\mathbf{Z}_{p^2}$ is equivalent to a code $\mathcal{C}$ with generator matrix

$$\begin{bmatrix} I_{k_1} & A \\ 0 & pB \end{bmatrix}$$

where $A \in M_{k_1 \times (n-k_1)}(\mathbf{Z})$, $B \in M_{k_2 \times (n-k_1)}(\mathbf{Z})$. Note that

$$\begin{bmatrix} I_{k_1} & A \end{bmatrix}$$

is a generator matrix of the residue code of $\mathcal{C}$, and

$$\begin{bmatrix} I_{k_1} & A \\ 0 & B \end{bmatrix}$$

is a generator matrix of the torsion code of $\mathcal{C}$. We say that the code $\mathcal{C}$ has type $\{k_1, k_2\}$.

The following lemma is due to Conway and Sloane [5, p. 34].

Lemma 2.1. *Let $\mathcal{C}$ be a self-orthogonal code of length n over $\mathbf{Z}_4$. Then $\pi(\mathcal{C})$ is doubly even and $\pi(\mathcal{C}) \subset \iota^{-1}(\mathcal{C}) \subset \pi(\mathcal{C})^\perp$ holds.*

We define the Euclidean weight in $\mathbf{Z}_4$ by $\text{wt}_e(0) = 0$, $\text{wt}_e(1) = \text{wt}_e(3) = 1$ and $\text{wt}_e(2) = 4$. The Euclidean weight of a vector $x = (x_1, \dots, x_n) \in \mathbf{Z}_4^n$ is defined by

$$\text{wt}_e(x) = \sum_{i=1}^n \text{wt}_e(x_i).$$

A quaternary code is said to be even if the Euclidean weight of every codeword is divisible by 8. Every quaternary even code is self-orthogonal. A quaternary self-orthogonal code with generator matrix all of whose row vectors have Euclidean weight divisible by 8 is even, by [3, Lemma 2.2]. If a quaternary even code contains a codeword all of whose coordinates are ± 1 , then the length is divisible by 8. A quaternary even self-dual code is also called a quaternary Type II code. It is known (see [7, Lemma 2.2]) that a quaternary Type II code contains a codeword all of whose coordinates are ± 1 , but in some earlier literature (see for example [4]), a quaternary Type II code is assumed to contain the all-ones vector $\mathbf{1}$.

3 Some mappings on matrices

Let K be a field, m a positive integer. We denote by $\text{Sym}_m(K)$ the set of symmetric $m \times m$ matrices and $\text{Alt}_m(K)$ the set of alternating $m \times m$ matrices over K . For a square matrix A of order n , we denote by $\text{diag}(A)$ the n -dimensional vector composed by the diagonal entries of A , and by

$\text{Diag}(A)$ the diagonal matrix whose diagonal entries are those of A . If the characteristic of K is 2, then

$$\text{Alt}_m(K) = \{A \in \text{Sym}_m(K) \mid \text{diag}(A) = 0\}.$$

Lemma 3.1. *If $A \in M_{m \times n}(K)$ has rank m , then the image of the map*

$$\begin{array}{ccc} \Psi_A : M_{m \times n}(K) & \rightarrow & M_m(K) \\ N & \mapsto & AN^t + NA^t \end{array}$$

is $\text{Sym}_m(K)$ if $\text{char } K \neq 2$ and $\text{Alt}_m(K)$ if $\text{char } K = 2$. Moreover, for $\text{char } K = 2$, the map

$$\begin{array}{ccc} \Phi_A : M_{m \times n}(K) & \rightarrow & \text{Sym}_m(K) \\ N & \mapsto & AN^t + NA^t + \text{Diag}(AN^t) \end{array}$$

is surjective.

Proof. Since $\text{rank } A = m$, $A(M_{n \times m}(K)) = M_m(K)$ holds. Thus we have

$$\begin{aligned} \Psi_A(M_{m \times n}(K)) &= \{AN^t + NA^t \mid N \in M_{m \times n}(K)\} \\ &= \{S + S^t \mid S \in M_m(K)\} \\ &= \begin{cases} \text{Sym}_m(K) & \text{if } \text{char } K \neq 2, \\ \text{Alt}_m(K) & \text{if } \text{char } K = 2. \end{cases} \end{aligned}$$

For $\text{char } K = 2$,

$$\begin{aligned} \text{Sym}_m(K) &= \{R + R^t + \text{Diag}(R) \mid R \in M_m(K) \text{ is upper triangular}\} \\ &\subset \{S + S^t + \text{Diag}(S) \mid S \in M_m(K)\} \\ &= \{AN^t + NA^t + \text{Diag}(AN^t) \mid N \in M_{m \times n}(K)\} \\ &= \Phi_A(M_{m \times n}(K)) \subset \text{Sym}_m(K). \end{aligned}$$

Thus $\Phi_A(M_{m \times n}(K)) = \text{Sym}_m(K)$. □

Lemma 3.2. *Let $\text{char } K = 2$, $\text{rank } A = m$ for $A \in M_{m \times n}(K)$ and the vector $\mathbf{1}$ does not belong to the row space of A . Define the map*

$$\begin{array}{ccc} \alpha : M_{m \times n}(K) & \rightarrow & K^m \\ N & \mapsto & \mathbf{1}N^t. \end{array}$$

Then the map

$$\begin{array}{ccc} \Phi_A \oplus \alpha : M_{m \times n}(K) & \rightarrow & \text{Sym}_m(K) \oplus K^m \\ N & \mapsto & (AN^t + NA^t + \text{Diag}(AN^t), \mathbf{1}N^t) \end{array}$$

is surjective.

Proof. Since Φ_A is surjective, it suffices to show that the restriction of α to $\text{Ker } \Phi_A$ is surjective. By the assumption,

$$\text{Ker } A \supsetneq \text{Ker } \begin{bmatrix} \mathbf{1} \\ A \end{bmatrix}.$$

This implies

$$K = \{\mathbf{1}x^t \mid x \in K^n, Ax^t = 0\},$$

and hence

$$\begin{aligned} \alpha(\text{Ker } \Phi_A) &\supseteq \{\mathbf{1}N^t \mid N \in M_{m \times n}(K), AN^t = 0\} \\ &= K^m. \end{aligned}$$

□

4 Codes over $\mathbf{Z}_{p^2}$ with prescribed residue and torsion

Throughout this section, we let p be a prime, and $\mathcal{C}_1, \mathcal{C}_2$ codes of length n over $\mathbf{Z}_p$ such that $\mathcal{C}_1$ has generator matrix

$$\begin{bmatrix} I & A \end{bmatrix}, \quad (2)$$

$\mathcal{C}_2$ has generator matrix

$$\begin{bmatrix} I & A \\ 0 & B \end{bmatrix}, \quad (3)$$

$A \in M_{k_1 \times (n-k_1)}(\mathbf{Z})$, $B \in M_{k_2 \times (n-k_1)}(\mathbf{Z})$ and $\dim \mathcal{C}_1 = k_1$, $\dim \mathcal{C}_2 = k_1 + k_2$.

A code $\mathcal{C}$ of length n over $\mathbf{Z}_{p^2}$ is free if $\mathcal{C}$ is isomorphic to the direct sum of copies of $\mathbf{Z}_{p^2}$, as a $\mathbf{Z}_{p^2}$ -module, or equivalently, $\pi(\mathcal{C}) = \iota^{-1}(\mathcal{C})$. We first compute the number of free self-orthogonal codes with given residue, then compute the number of self-orthogonal codes with given residue and torsion.

Lemma 4.1. *If $\mathcal{C}$ is a code of length n over $\mathbf{Z}_{p^2}$ satisfying $\pi(\mathcal{C}) = \mathcal{C}_1$ and $\iota^{-1}(\mathcal{C}) = \mathcal{C}_2$, then there exists a matrix $N \in M_{k_1 \times (n-k_1)}(\mathbf{Z})$ such that*

$$\begin{bmatrix} I & A + pN \\ 0 & pB \end{bmatrix} \quad (4)$$

is a generator matrix of $\mathcal{C}$. Moreover, if $k_2 = 0$, that is, $\pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1$, such a matrix N is unique modulo p .

Proof. Since $\pi(\mathcal{C}) = \mathcal{C}_1$ and $\iota^{-1}(\mathcal{C}) = \mathcal{C}_2$,

$$\mathcal{C} \supset \mathbf{Z}_{p^2}^{k_1+k_2} \begin{bmatrix} I + pM_1 & A + pM_2 \\ 0 & pB \end{bmatrix}$$

for some $M_1 \in M_{k_1}(\mathbf{Z})$, $M_2 \in M_{k_1 \times (n-k_1)}(\mathbf{Z})$. Then,

$$\begin{aligned} \mathcal{C} &\supset \mathbf{Z}_{p^2}^{k_1+k_2} \begin{bmatrix} I - pM_1 & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} I + pM_1 & A + pM_2 \\ 0 & pB \end{bmatrix} \\ &= \mathbf{Z}_{p^2}^{k_1+k_2} \begin{bmatrix} I & A + p(M_2 - M_1A) \\ 0 & pB \end{bmatrix}. \end{aligned}$$

Taking $N = M_2 - M_1A$, we have

$$\begin{aligned} |\mathcal{C}| &\geq \left| \mathbf{Z}_{p^2}^{k_1+k_2} \begin{bmatrix} I & A + pN \\ 0 & pB \end{bmatrix} \right| \\ &= p^{2k_1+k_2} \\ &= |\mathcal{C}_2| |\mathcal{C}_1| \\ &= |\mathcal{C}| \end{aligned}$$

by (1). Thus, $\mathcal{C}$ has a generator matrix (4). It remains to show that for $k_2 = 0$, the matrix N in (4) is determined uniquely by $\mathcal{C}$. Indeed, if $N_1, N_2 \in M_{k_1 \times (n-k_1)}(\mathbf{Z})$ and

$$\mathbf{Z}_{p^2}^{k_1} \begin{bmatrix} I & A + pN_1 \end{bmatrix} = \mathbf{Z}_{p^2}^{k_1} \begin{bmatrix} I & A + pN_2 \end{bmatrix},$$

then $A + pN_1 \equiv A + pN_2 \pmod{p^2}$, and hence $N_1 \equiv N_2 \pmod{p}$. $\square$

For the remainder of this section, we assume $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_1^\perp$. This implies

$$I + AA^t \equiv 0 \pmod{p} \tag{5}$$

and

$$AB^t \equiv 0 \pmod{p}. \tag{6}$$

Moreover, when $p = 2$, we further assume $\mathcal{C}_1$ to be doubly even, or equivalently,

$$\text{diag}(I + AA^t) \equiv 0 \pmod{4}. \tag{7}$$

This is justified by Lemma 2.1.

Lemma 4.2. *The number of free self-orthogonal codes $\mathcal{C} \subseteq \mathbf{Z}_{p^2}^n$ such that $\pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1$ is $p^{k_1(2n-3k_1-1)/2}$ for odd primes p and $2^{k_1(2n-3k_1+1)/2}$ for $p = 2$.*

Proof. By Lemma 4.1, a code $\mathcal{C} \subset \mathbf{Z}_{p^2}^n$ such that $\pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1$ has a generator matrix $[I \ A + pN]$ for some matrix $N \in M_{k_1 \times (n-k_1)}(\mathbf{Z})$ which is unique modulo p . Observe that $\mathcal{C}$ is self-orthogonal if and only if

$$I + AA^t + p(AN^t + NA^t) \equiv 0 \pmod{p^2}. \quad (8)$$

By (5), $-\frac{1}{p}(I + AA^t) \pmod{p}$ is a symmetric matrix over $\mathbf{Z}_p$ if p is odd, and by (7), it is an alternating matrix over $\mathbf{Z}_p$ if $p = 2$. Therefore by Lemma 3.1,

$$\begin{aligned} & |\{\mathcal{C} \mid \mathcal{C} \text{ is self-orthogonal, } \pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1\}| \\ &= |\{N \pmod{p} \mid N \in M_{k_1 \times (n-k_1)}(\mathbf{Z}) \text{ satisfies (8)}\}| \\ &= \left| \left\{ N \mid N \in M_{k_1 \times (n-k_1)}(\mathbf{Z}_p) \text{ and } \Psi_A(N) = -\frac{1}{p}(I + AA^t) \pmod{p} \right\} \right| \\ &= \left| \Psi_A^{-1} \left(-\frac{1}{p}(I + AA^t) \pmod{p} \right) \right| \\ &= |\text{Ker } \Psi_A| \\ &= \begin{cases} \frac{|M_{k_1 \times (n-k_1)}(\mathbf{Z}_p)|}{|\text{Sym}_{k_1}(\mathbf{Z}_p)|} & \text{if } p \text{ is odd,} \\ \frac{|M_{k_1 \times (n-k_1)}(\mathbf{Z}_2)|}{|\text{Alt}_{k_1}(\mathbf{Z}_2)|} & \text{if } p = 2, \end{cases} \\ &= \begin{cases} p^{k_1(n-k_1)-k_1(k_1+1)/2} & \text{if } p \text{ is odd,} \\ 2^{k_1(n-k_1)-k_1(k_1-1)/2} & \text{if } p = 2. \end{cases} \end{aligned}$$

□

Let us consider sets

$$\begin{aligned} X &= \{\mathcal{C} \mid \mathcal{C} \text{ is self-orthogonal, } \pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1\} \text{ and} \\ X' &= \{\mathcal{C}' \mid \mathcal{C}' \text{ is self-orthogonal, } \pi(\mathcal{C}') = \mathcal{C}_1, \iota^{-1}(\mathcal{C}') = \mathcal{C}_2\}. \end{aligned}$$

By Lemma 4.2, we have $|X| = p^{k_1(2n-3k_1-1)/2}$ if p is odd and $|X| = 2^{k_1(2n-3k_1+1)/2}$ if $p = 2$.

Lemma 4.3. *If $\mathcal{C} \in X$, then there exists a unique $\mathcal{C}' \in X'$ containing $\mathcal{C}$.*

Proof. By Lemma 4.1, $\mathcal{C}$ has a generator matrix $\begin{bmatrix} I & A + pN \end{bmatrix}$ for some N . Then the code $\mathcal{C}'_0$ with generator matrix

$$\begin{bmatrix} I & A + pN \\ 0 & pB \end{bmatrix}$$

satisfies $\pi(\mathcal{C}'_0) = \mathcal{C}_1$ and $\iota^{-1}(\mathcal{C}'_0) = \mathcal{C}_2$. Since $\mathcal{C} \in X$, (6) implies that $\mathcal{C}'_0$ is self-orthogonal, hence $\mathcal{C}'_0 \in X'$. If $\mathcal{C}' \in X'$ and $\mathcal{C} \subset \mathcal{C}'$, then Lemma 4.1 implies $\mathbf{Z}_{p^2}^{k_2} \begin{bmatrix} I & pB \end{bmatrix} \subset \mathcal{C}'$. This, together with $\mathcal{C} \subset \mathcal{C}'$ forces $\mathcal{C}'_0 \subset \mathcal{C}'$, and hence $\mathcal{C}'_0 = \mathcal{C}'$. $\square$

Lemma 4.4. *Let $\mathcal{C}' \in X'$. Then $|\{\mathcal{C} \in X \mid \mathcal{C} \subset \mathcal{C}'\}| = p^{k_1 k_2}$.*

Proof. Let $\begin{bmatrix} I & A + pN \\ 0 & pB \end{bmatrix}$ be a generator matrix of $\mathcal{C}'$. Consider the map

$$\varphi : M_{k_1 \times k_2}(\mathbf{Z}_p) \rightarrow \{\mathcal{C} \in X \mid \mathcal{C} \subset \mathcal{C}'\}$$

$$M \bmod p \mapsto \mathbf{Z}_{p^2}^{k_1} \begin{bmatrix} I & A + p(N + MB) \end{bmatrix}.$$

We claim that this map is bijective. Suppose $M_1, M_2 \in M_{k_1 \times k_2}(\mathbf{Z})$, $\varphi(M_1) = \varphi(M_2)$. Then we have $A + p(N + M_1 B) \equiv A + p(N + M_2 B) \pmod{p^2}$. Since $\text{rank}(B \bmod p) = k_2$, we conclude $M_1 \equiv M_2 \pmod{p}$, which shows that φ is injective. Next we show that φ is surjective. Suppose $\mathcal{C} \in X$ and $\mathcal{C} \subset \mathcal{C}'$. Then by Lemma 4.1, $\mathcal{C} = \mathbf{Z}_{p^2}^{k_1} \begin{bmatrix} I & A + pF \end{bmatrix}$ for some matrix F . Since $\mathcal{C} \subset \mathcal{C}'$, $A + pF \equiv A + pN + pMB \pmod{p^2}$ for some matrix M . Then we have $F \equiv N + MB \pmod{p}$, so we conclude that φ is surjective. Therefore, the map φ is bijective. It follows that the number of codes $\mathcal{C} \in X$ contained in $\mathcal{C}'$ is $p^{k_1 k_2}$. $\square$

Theorem 4.5. *Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be codes of length n over $\mathbf{Z}_p$ where $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_1^\perp$. Assume further that $\mathcal{C}_1$ is doubly even when $p = 2$. If $\dim \mathcal{C}_1 = k_1$ and $\dim \mathcal{C}_2 = k_1 + k_2$, then the number of self-orthogonal codes $\mathcal{C}$ of length n over $\mathbf{Z}_{p^2}$ such that $\pi(\mathcal{C}) = \mathcal{C}_1$ and $\iota^{-1}(\mathcal{C}) = \mathcal{C}_2$ is $p^{k_1(2n-3k_1-1-2k_2)/2}$ for odd primes p and $2^{k_1(2n-3k_1+1-2k_2)/2}$ for $p = 2$.*

Proof. We may assume without loss of generality that $\mathcal{C}_1$ and $\mathcal{C}_2$ are codes with generator matrices given by (2) and (3), respectively. By Lemma 4.3

and Lemma 4.4, we have

$$\begin{aligned}
|X'| &= p^{-k_1 k_2} \sum_{\mathcal{C}' \in X'} |\{\mathcal{C} \in X \mid \mathcal{C} \subset \mathcal{C}'\}| \\
&= p^{-k_1 k_2} \sum_{\mathcal{C} \in X} |\{\mathcal{C}' \in X' \mid \mathcal{C} \subset \mathcal{C}'\}| \\
&= p^{-k_1 k_2} |X|.
\end{aligned}$$

The result then follows from Lemma 4.2. $\square$

5 Quaternary even codes with prescribed residue and torsion

We are mainly concerned with the enumeration of codes containing the vector $\mathbf{1}$, or codes containing a vector each of whose coordinate is 1 or -1 . This restriction forces the length of a code to be a multiple of 8. First, let us consider such codes containing $\mathbf{1}$. Let $n \in 8\mathbf{Z}$, $1 \leq k_1 \leq \frac{n}{2}$, $\mathcal{C}_1, \mathcal{C}_2$ binary codes of length n such that $\mathcal{C}_1$ is doubly even and has generator matrix

$$\begin{bmatrix} 1 & \mathbf{1} & \mathbf{1} \\ 0 & I_{k_1-1} & A \end{bmatrix}, \quad (9)$$

$\mathcal{C}_2$ has generator matrix

$$\begin{bmatrix} 1 & \mathbf{1} & \mathbf{1} \\ 0 & I_{k_1-1} & A \\ 0 & 0 & B \end{bmatrix}, \quad (10)$$

$A \in M_{(k_1-1) \times (n-k_1)}(\mathbf{Z})$, $B \in M_{k_2 \times (n-k_1)}(\mathbf{Z})$ and $\dim \mathcal{C}_1 = k_1$, $\dim \mathcal{C}_2 = k_1 + k_2$. Moreover, we assume that $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_1^\perp$ and $\mathcal{C}_1$ is doubly even. Then the matrices A and B satisfy (5)–(7), and

$$\mathbf{1}B^t \equiv 0 \pmod{2}. \quad (11)$$

We may assume without loss of generality that the entries of the matrix A are all 0 or 1. Then

$$\mathbf{1} + \mathbf{1}A^t \equiv 0 \pmod{4}. \quad (12)$$

Lemma 5.1. *The vector $\mathbf{1}$ does not belong to the row space of the matrix A over $\mathbf{Z}_2$.*

Proof. If $k_1 = 1$, the assertion is trivial, so assume $k_1 > 1$. Suppose that $\mathbf{1}$ is in the row space of A over $\mathbf{Z}_2$. Then $bA \equiv \mathbf{1} \pmod{2}$ for some vector $b \in \mathbf{Z}^{k_1-1}$. By (5) and (12), we have

$$\begin{aligned} 0 &\equiv [b(\mathbf{1} + \mathbf{1}A^t)^t \quad b(I + AA^t)] \pmod{2} \\ &\equiv [b\mathbf{1}^t + \mathbf{1}\mathbf{1}^t \quad b + \mathbf{1}A^t] \pmod{2} \\ &\equiv [b\mathbf{1}^t + (n - k_1) \quad b + \mathbf{1}] \pmod{2} \quad (\text{by (12)}). \end{aligned}$$

Therefore, $b \equiv \mathbf{1} \pmod{2}$ and

$$b\mathbf{1}^t + (n - k_1) \equiv (k_1 - 1) + (n - k_1) \not\equiv 0 \pmod{2}.$$

This is a contradiction. □

Lemma 5.2. *Define the maps*

$$\begin{aligned} \alpha : M_{(k_1-1) \times (n-k_1)}(\mathbf{Z}_2) &\rightarrow \mathbf{Z}_2^{k_1-1} \\ N \bmod 2 &\mapsto \mathbf{1}N^t \bmod 2 \end{aligned}$$

and

$$\begin{aligned} \Phi_A : M_{(k_1-1) \times (n-k_1)}(\mathbf{Z}_2) &\rightarrow \text{Sym}_{k_1-1}(\mathbf{Z}_2) \\ N \bmod 2 &\mapsto AN^t + NA^t + \text{Diag}(AN^t) \bmod 2. \end{aligned}$$

For $N \in M_{(k_1-1) \times (n-k_1)}(\mathbf{Z})$, let

$$\mathcal{C} = \mathbf{Z}_4^{k_1} \begin{bmatrix} 1 & \mathbf{1} & \mathbf{1} \\ 0 & I_{k_1-1} & A + 2N \end{bmatrix}.$$

Then $\mathcal{C}$ is an even code if and only if $N \bmod 2$ belongs to the set

$$\text{Ker } \alpha \cap \Phi_A^{-1} \left(\frac{1}{2}(I + AA^t) + \frac{1}{4}(I + \text{Diag } AA^t) \bmod 2 \right).$$

Proof. $\mathcal{C}$ is self-orthogonal if and only if

$$\mathbf{1} + \mathbf{1}A^t + 2\mathbf{1}N^t \equiv 0 \pmod{4}, \quad (13)$$

$$I + AA^t + 2(AN^t + NA^t) \equiv 0 \pmod{4}. \quad (14)$$

In view of (12), (13) is equivalent to $N \bmod 2 \in \text{Ker } \alpha$. Moreover, when $\mathcal{C}$ is self-orthogonal, $\mathcal{C}$ is even if and only if

$$\text{diag}(I + AA^t + 2(AN^t + NA^t) + 4NN^t) \equiv 0 \pmod{8}. \quad (15)$$

Since $\text{diag } NN^t \equiv \mathbf{1}N^t \equiv 0 \pmod{2}$, this is equivalent to

$$\text{Diag}(AN^t) \equiv \frac{1}{4}(I + \text{Diag } AA^t) \pmod{2}. \quad (16)$$

Therefore, the code $\mathcal{C}$ is even if and only if $N \bmod 2 \in \text{Ker } \alpha$ and conditions (14) and (16) are satisfied. The latter two conditions are equivalent to

$$\Phi_A(N \bmod 2) = \frac{1}{2}(I + AA^t) + \frac{1}{4}(I + \text{Diag } AA^t) \bmod 2.$$

□

Lemma 5.3. *The number of free quaternary even codes $\mathcal{C}$ of length n containing $\mathbf{1}$ such that $\pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1$ is $2^{(k_1-1)(2n-3k_1-2)/2}$.*

Proof. By Lemma 5.2, we have

$$\begin{aligned} & |\{\mathcal{C} \mid \mathcal{C} \text{ is even, } \pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1, \mathbf{1} \in \mathcal{C}\}| \\ &= \left| \text{Ker } \alpha \cap \Phi_A^{-1} \left(\frac{1}{2}(I + AA^t) + \frac{1}{4}(I + \text{Diag } AA^t) \bmod 2 \right) \right| \\ &= \left| (\Phi_A \oplus \alpha)^{-1} \left(\left(\frac{1}{2}(I + AA^t) + \frac{1}{4}(I + \text{Diag } AA^t) \bmod 2 \right), 0 \right) \right|. \end{aligned}$$

By Lemma 5.1, $\mathbf{1}$ does not belong to the row space of A over $\mathbf{Z}_2$. Thus, the map $\Phi_A \oplus \alpha$ is surjective by Lemma 3.2. Therefore, the number of even codes $\mathcal{C}$ containing $\mathbf{1}$ such that $\pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1$ is

$$|\text{Ker}(\Phi_A \oplus \alpha)| = 2^{(k_1-1)(n-k_1)-(k_1-1)-k_1(k_1-1)/2}.$$

□

Let us consider the sets

$$\begin{aligned} Y &= \{\mathcal{C} \mid \mathcal{C} \text{ is quaternary even, } \pi(\mathcal{C}) = \iota^{-1}(\mathcal{C}) = \mathcal{C}_1, \mathbf{1} \in \mathcal{C}\} \text{ and} \\ Y' &= \{\mathcal{C}' \mid \mathcal{C}' \text{ is quaternary even, } \pi(\mathcal{C}') = \mathcal{C}_1, \iota^{-1}(\mathcal{C}') = \mathcal{C}_2, \mathbf{1} \in \mathcal{C}'\}. \end{aligned}$$

By Lemma 5.3, $|Y| = 2^{(k_1-1)(2n-3k_1-2)/2}$.

Lemma 5.4. *If $\mathcal{C} \in Y$, then there exists a unique $\mathcal{C}' \in Y'$ containing $\mathcal{C}$.*

Proof. The unique code $\mathcal{C}'$ given in Lemma 4.3 is even. The matrix $B \bmod 2$ has even weight rows by (11) so $2B \bmod 4$ has rows with Euclidean weights divisible by 8. $\square$

Lemma 5.5. *Let $\mathcal{C}' \in Y'$. Then $|\{\mathcal{C} \in Y \mid \mathcal{C} \subset \mathcal{C}'\}| = 2^{(k_1-1)k_2}$.*

Proof. Let $\begin{bmatrix} 1 & \mathbf{1} & \mathbf{1} \\ 0 & I_{k_1-1} & A + 2N \\ 0 & 0 & 2B \end{bmatrix}$ be a generator matrix of $\mathcal{C}'$. Consider the map

$$\varphi' : M_{(k_1-1) \times k_2}(\mathbf{Z}_2) \rightarrow \{\mathcal{C} \in Y \mid \mathcal{C} \subset \mathcal{C}'\}$$

$$M \bmod 2 \mapsto \mathbf{Z}_4^{k_1} \begin{bmatrix} 1 & \mathbf{1} & \mathbf{1} \\ 0 & I_{k_1-1} & A + 2(N + MB) \end{bmatrix}.$$

We claim that this map is bijective. Suppose $M_1, M_2 \in M_{(k_1-1) \times k_2}(\mathbf{Z})$ and $\varphi'(M_1) = \varphi'(M_2)$. Using same approach as in the proof of Lemma 4.4, we can show that the map φ' is well-defined and injective with $M_1 \equiv M_2 \pmod{2}$. Next we show that the map is surjective. Suppose $\mathcal{C} \in Y$ and $\mathcal{C} \subset \mathcal{C}'$, then $\mathcal{C} = \mathbf{Z}_4^{k_1} \begin{bmatrix} 1 & \mathbf{1} & \mathbf{1} \\ 0 & I_{k_1-1} & A + 2F \end{bmatrix}$ for some matrix F . Since $\mathcal{C} \subset \mathcal{C}'$, $A + 2F \equiv A + 2N + 2MB \pmod{4}$ for some matrix M . Then we have $F \equiv N + MB \pmod{2}$, so we conclude that φ' is surjective. Hence, the map φ' is bijective. Therefore the number of codes $\mathcal{C} \in Y$ contained in $\mathcal{C}'$ is $2^{(k_1-1)k_2}$. $\square$

Theorem 5.6. *Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be binary codes of length n where $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_1^\perp$. If $\mathcal{C}_1$ is doubly even, $\mathbf{1} \in \mathcal{C}_1$ and $\dim \mathcal{C}_1 = k_1$ and $\dim \mathcal{C}_2 = k_1 + k_2$, then the number of quaternary even codes $\mathcal{C}$ containing $\mathbf{1}$ such that $\pi(\mathcal{C}) = \mathcal{C}_1$ and $\iota^{-1}(\mathcal{C}) = \mathcal{C}_2$ is $2^{(k_1-1)(2n-3k_1-2-2k_2)/2}$.*

Proof. We may assume without loss of generality that $\mathcal{C}_1$ and $\mathcal{C}_2$ are codes with generator matrices given by (9) and (10), respectively. By Lemma 5.4 and Lemma 5.5, we have

$$\begin{aligned} |Y'| &= 2^{-(k_1-1)k_2} \sum_{\mathcal{C}' \in Y'} |\{\mathcal{C} \in Y \mid \mathcal{C} \subset \mathcal{C}'\}| \\ &= 2^{-(k_1-1)k_2} \sum_{\mathcal{C} \in Y} |\{\mathcal{C}' \in Y' \mid \mathcal{C} \subset \mathcal{C}'\}| \\ &= 2^{-(k_1-1)k_2} |Y|. \end{aligned}$$

The result then follows from Lemma 5.3. $\square$

Theorem 5.7. *Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be binary codes of length n where $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_1^\perp$. If $\mathcal{C}_1$ is doubly even, $\mathbf{1} \in \mathcal{C}_1$, $\dim \mathcal{C}_1 = k_1$ and $\dim \mathcal{C}_2 = k_1 + k_2$, then the number of quaternary even codes $\mathcal{C}$ containing an element of $\{\pm 1\}^n$ such that $\pi(\mathcal{C}) = \mathcal{C}_1$ and $\iota^{-1}(\mathcal{C}) = \mathcal{C}_2$ is $2^{(n-k_1-k_2)+(k_1-1)(2n-3k_1-2-2k_2)/2}$.*

Proof. Consider the sets

$$Z = \{\pm 1\}^n \text{ and}$$

$$U = \{\mathcal{C} \mid \mathcal{C} \text{ is quaternary even, } \pi(\mathcal{C}) = \mathcal{C}_1, \iota^{-1}(\mathcal{C}) = \mathcal{C}_2, Z \cap \mathcal{C} \neq \emptyset\}.$$

If $\mathcal{C} \in U$, then there exists $z \in Z \cap \mathcal{C}$. Then $Z \cap \mathcal{C} = \{z + \iota(x) \mid x \in \mathcal{C}_2\}$, hence $|Z \cap \mathcal{C}| = |\mathcal{C}_2|$. Therefore, we have

$$\begin{aligned} 2^{k_1+k_2}|U| &= |\mathcal{C}_2||U| \\ &= \sum_{\mathcal{C} \in U} |Z \cap \mathcal{C}| \\ &= \sum_{z \in Z} |\{\mathcal{C} \in U \mid z \in \mathcal{C}\}| \\ &= \sum_{z \in Z} |\{\mathcal{C} \in U \mid \mathbf{1} \in \mathcal{C}\}| \\ &= |Z| 2^{(k_1-1)(2n-3k_1-2-2k_2)/2} \quad (\text{by Theorem 5.6}) \\ &= 2^{n+(k_1-1)(2n-3k_1-2-2k_2)/2}. \end{aligned}$$

$\square$

6 Main Results

Let $\sigma(n, k_1)$ denote the number of distinct doubly even binary codes of length n and dimension k_1 , and let $\sigma_p(n, k_1)$ be the number of distinct self-orthogonal p -ary codes, with p an odd prime, of length n and dimension k_1 . Let $\sigma_1(n, k_1)$ be the number of distinct doubly even binary codes of length n and dimension k_1 containing $\mathbf{1}$. The value of $\sigma_p(n, k_1)$ is given in [8, 9]. The values of $\sigma(n, k_1)$ and $\sigma_1(n, k_1)$ are given in [6]. All of these values can also be derived from

[11]. For $k \leq n$, we define the Gaussian coefficient $\begin{bmatrix} n \\ k \end{bmatrix}_p$ as

$$\begin{bmatrix} n \\ k \end{bmatrix}_p = \frac{(p^n - 1)(p^n - p) \cdots (p^n - p^{k-1})}{(p^k - 1)(p^k - p) \cdots (p^k - p^{k-1})}.$$

Corollary 1. *The number of distinct self-orthogonal codes of length n over $\mathbf{Z}_{p^2}$ of type $\{k_1, k_2\}$ is*

$$M_{p^2}(k_1, k_2) = \sigma_p(n, k_1) \begin{bmatrix} n - 2k_1 \\ k_2 \end{bmatrix}_p p^{k_1(2n-3k_1-1-2k_2)/2}, \quad (17)$$

for odd primes p , and

$$M_4(k_1, k_2) = \sigma(n, k_1) \begin{bmatrix} n - 2k_1 \\ k_2 \end{bmatrix}_2 2^{k_1(2n-3k_1+1-2k_2)/2}, \quad (18)$$

for $p = 2$.

Proof. Given a self-orthogonal $[n, k_1]$ code $\mathcal{C}_1$, we have $\begin{bmatrix} n - 2k_1 \\ k_2 \end{bmatrix}_p$ codes $\mathcal{C}_2$ such that $\mathcal{C}_1 \subseteq \mathcal{C}_2 \subseteq \mathcal{C}_1^\perp$. The result follows from Theorem 4.5. $\square$

As an example, we consider the case $n = 4$, $k_1 = k_2 = 1$, and $p = 3$. Let $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{C}_4$ be the self-orthogonal codes over $\mathbf{Z}_9$ of type $\{1, 1\}$ with generator matrices

$$\begin{bmatrix} 1 & 1 & 4 & 0 \\ 0 & 3 & 6 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 4 & 3 \\ 0 & 3 & 6 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 4 & 6 \\ 0 & 3 & 6 & 3 \end{bmatrix}, \begin{bmatrix} 1 & 7 & 7 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix},$$

respectively. The orders of their automorphism groups are 24, 12, 4 and 8, respectively. Thus, we have

$$\sum_{i=1}^4 \frac{|E|}{|\text{Aut } \mathcal{C}_i|} = \frac{2^4 4!}{24} + \frac{2^4 4!}{12} + \frac{2^4 4!}{4} + \frac{2^4 4!}{8} = 16 + 32 + 96 + 48 = 192.$$

Using Corollary 1, we have

$$\begin{aligned} M_9(1, 1) &= \sigma_3(4, 1) \begin{bmatrix} 4 - 2 \\ 1 \end{bmatrix}_3 3^{3-1-1} \\ &= 16 \cdot 4 \cdot 3 = 192. \end{aligned}$$

This implies that $\{\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{C}_4\}$ is a complete set of representatives for equivalence classes of self-orthogonal codes of length 4 and type $\{1, 1\}$ over $\mathbf{Z}_9$.

As a consequence of Corollary 1, we have

Corollary 2. *The number of distinct self-dual codes over $\mathbf{Z}_{p^2}$ of length n is*

$$\sum_{0 \leq k_1 \leq \lfloor \frac{n}{2} \rfloor} M_{p^2}(k_1, n - 2k_1). \quad (19)$$

Corollary 2 agrees with the results in [6], [1], and [2].

Corollary 3. *The number of distinct quaternary even codes of length n containing $\mathbf{1}$, of type $\{k_1, k_2\}$ is*

$$D_1(k_1, k_2) = \sigma_1(n, k_1) \begin{bmatrix} n - 2k_1 \\ k_2 \end{bmatrix}_2 2^{(k_1-1)(2n-3k_1-2-2k_2)/2}, \quad (20)$$

and the number of distinct quaternary even codes of length n containing an element of $\{\pm 1\}^n$, of type $\{k_1, k_2\}$ is

$$D(k_1, k_2) = \sigma_1(n, k_1) \begin{bmatrix} n - 2k_1 \\ k_2 \end{bmatrix}_2 2^{(n-k_1-k_2)+(k_1-1)(2n-3k_1-2-2k_2)/2}. \quad (21)$$

Proof. Given a binary doubly even $[n, k_1]$ code $\mathcal{C}_1$ containing $\mathbf{1}$, we have $\begin{bmatrix} n - 2k_1 \\ k_2 \end{bmatrix}_2$ codes $\mathcal{C}_2$ such that $\mathcal{C}_1 \subseteq \mathcal{C}_2 \subseteq \mathcal{C}_1^\perp$. Thus, (20) follows from Theorem 5.6. Similarly, (21) follows from Theorem 5.7. $\square$

And as a consequence of Corollary 3, we have

Corollary 4. *The number of distinct quaternary Type II codes of length n containing $\mathbf{1}$ is*

$$\sum_{0 \leq k_1 \leq \frac{n}{2}} D_1(k_1, n - 2k_1), \quad (22)$$

and the number of distinct quaternary Type II codes of length n containing an element of $\{\pm 1\}^n$ is

$$\sum_{0 \leq k_1 \leq \frac{n}{2}} D(k_1, n - 2k_1). \quad (23)$$

The formula (23) agrees with [6, Theorem 6].

References

- [1] J.M.P. Balmaceda, R.A.L. Betty and F.R. Nemenzo, On the number of distinct self-dual codes over $\mathbf{Z}_9$, *Matimyas Matematika*, 26 (2003), 9–17.
- [2] J.M.P. Balmaceda, R.A.L. Betty and F.R. Nemenzo, Mass formula for self-dual codes over $\mathbf{Z}_{p^2}$, *Discrete Math.*, 308 (2008), 2984–3002 .
- [3] E. Bannai, S.T. Dougherty, M. Harada and M. Oura, Type II codes, even unimodular lattices and invariant rings, *IEEE Trans. Inform. Theory*, 45 (1999), 257–269.
- [4] A. Bonnetcaze, P. Solé, C. Bachoc and B. Mourrain, Type II codes over $\mathbf{Z}_4$, *IEEE Trans. Inform. Theory*, 43 (1997), 969–976.
- [5] J.H. Conway and N.J.A. Sloane, Self-dual codes over the integers modulo 4, *J. Combin. Theory Ser. A*, 62 (1993), 30–45.
- [6] P. Gaborit, Mass formulas for self-dual codes over $\mathbf{Z}_4$ and $\mathbf{F}_q + u\mathbf{F}_q$ rings, *IEEE Trans. Inform. Theory*, 42 (1996), 1222–1228.
- [7] M. Harada, P. Solé and P. Gaborit, Self-dual codes over $\mathbf{Z}_4$ and unimodular lattices: a survey, *Algebras and combinatorics (Hong Kong, 1997)*, 255–275, Springer, Singapore, 1999.
- [8] V.S. Pless, The number of isotropic subspaces in a finite geometry, *Atti. Accad. Naz. Lincei Rendic*, 39 (1965), 418–421.
- [9] V.S. Pless, On the uniqueness of the Golay codes, *J. Combin. Theory*, 5 (1968), 215–228.
- [10] E.M. Rains and N.J.A. Sloane, Self-Dual Codes, in “Handbook of Coding Theory,” V.S. Pless and W.C. Huffman, eds., Elsevier, Amsterdam, (1998), pp. 177–294.
- [11] D.K. Ray-Chaudhuri, Some results on quadrics in finite projective geometry based on Galois fields, *Canad. J. Math.*, 14 (1962), 129–138.